



## Cleaved-MMP-17 (Q129) rabbit pAb

**Cat#: orb763921 (Manual)** 

For research use only. Not intended for diagnostic use.

Product Name Cleaved-MMP-17 (Q129) rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human MMP17. AÅ range:110-159

Specificity Cleaved-MMP-17 (Q129) Polyclonal Antibody detects endogenous levels of

fragment of activated MMP-17 protein resulting from cleavage adjacent to

Q129.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Matrix metalloproteinase-17

Gene Name MMP17

Cellular localization [Isoform Long]: Cell membrane; Lipid-anchor, GPI-anchor; Extracellular

side. Secreted, extracellular space, extracellular matrix.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal





Concentration 1 mg/ml

**Observed band** 53kD

**Human Gene ID** 4326

**Human Swiss-Prot Number** Q9ULZ9

MMP17; MT4MMP; Matrix metalloproteinase-17; MMP-17; Membrane-type matrix metalloproteinase 4; MT-MMP 4; MTMMP4; Membrane-type-4 matrix metalloproteinase; MT4-MMP; MT4MMP **Alternative Names** 


**Background** This gene encodes a member of the peptidase M10 family and membranetype subfamily of matrix metalloproteinases (MMPs). Proteins in this family

are involved in the breakdown of extracellular matrix in normal

physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Members of this subfamily contain a transmembrane domain suggesting that these proteins are expressed at the cell surface rather than secreted. The encoded preproprotein is proteolytically processed to generate the mature protease. This protein is unique among the membrane-type matrix metalloproteinases in that it is anchored to the cell membrane via a glycosylphosphatidylinositol (GPI) anchor. Elevated expression of the

encoded protein has been observed in osteoarthritis and multiple human

cancers. [provided by RefSeq, Jan 2016],




Western Blot analysis of various cells using Cleaved-MMP-17 (Q129) Polyclonal Antibody





Explore. Bioreagents.



Western blot analysis of lysates from A549 cells, treated with etoposide 25uM 1h, using MMP17 (Cleaved-Gln129) Antibody. The lane on the right is blocked with the synthesized peptide.