



Bad (phospho Ser136) rabbit pAb

Cat#: orb764145 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Bad (phospho Ser136) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA:

1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human BAD around the phosphorylation site of Ser 136. AA range: 102-151

Specificity Phospho-Bad (S136) Polyclonal Antibody detects endogenous levels of Bad

protein only when phosphorylated at S136.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Bcl2 antagonist of cell death

Gene Name BAD

Cellular localization Mitochondrion outer membrane. Cytoplasm . Colocalizes with HIF3A in the

cytoplasm (By similarity). Upon phosphorylation, locates to the cytoplasm. .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Explore. Bioreagents.

1 mg/mlConcentration

Observed band

Human Gene ID 572

Human Swiss-Prot Number Q92934

BAD; BBC6; BCL2L8; Bcl2 antagonist of cell death; BAD; Bcl-2-binding component 6; Bcl-2-like protein 8; Bcl2-L-8; Bcl-XL/Bcl-2-associated death **Alternative Names**


promoter

The protein encoded by this gene is a member of the BCL-2 family. BCL-2 **Background**


family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL and BCL-2, and reversing their death repressor activity.


Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which

encode the same isoform. [provided by RefSeq, Jul 2008],



Western Blot analysis of various cells using Phospho-Bad (\$136) Polyclonal Antibody





Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using BAD (Phospho-Šer136) Antibody. The picture on the right is blocked with the phospho peptide.



