



Bcl-2 (phospho Thr56) rabbit pAb

Cat#: orb764148 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Bcl-2 (phospho Thr56) rabbit pAb

Host species Rabbit

Applications WB;IHC;IP;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunoprecipitation: 2-5 ug/mg lysate. Immunofluorescence: 1/200 -**Recommended dilutions**

1/1000. ELISA: 1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human BCL-2 around the phosphorylation site of Thr56. AA range:26-75

Phospho-Bcl-2 (T56) Polyclonal Antibody detects endogenous levels of Bcl-**Specificity**

2 protein only when phosphorylated at T56.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage**

Protein Name Apoptosis regulator Bcl-2

Gene Name BCL2

Cellular localization Mitochondrion outer membrane; Single-pass membrane protein. Nucleus

membrane ; Single-pass membrane protein . Endoplasmic reticulum membrane ; Single-pass membrane protein . Cytoplasm .

The antibody was affinity-purified from rabbit antiserum by affinity-**Purification**

chromatography using epitope-specific immunogen.





Clonality Polyclonal

Concentration 1 mg/ml

Observed band 26kD

596 **Human Gene ID**

Human Swiss-Prot Number P10415

Alternative Names BCL2; Apoptosis regulator Bcl-2

Background

BCL2, apoptosis regulator(BCL2) Homo sapiens This gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016],