



## c-Kit (phospho Tyr721) rabbit pAb

Cat#: orb764164 (Manual)

For research use only. Not intended for diagnostic use.

**Product Name** c-Kit (phospho Tyr721) rabbit pAb

**Host species** Rabbit

**Applications** WB;ELISA

**Species Cross-Reactivity** Human; Mouse; Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

**Immunogen** The antiserum was produced against synthesized peptide derived from

human c-Kit around the phosphorylation site of Tyr721. AA range:688-737

Phospho-c-Kit (Y721) Polyclonal Antibody detects endogenous levels of c-**Specificity** 

Kit protein only when phosphorylated at Y721.

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage** 

**Protein Name** Mast/stem cell growth factor receptor Kit

Gene Name KIT

Cellular localization

[Isoform 1]: Cell membrane; Single-pass type I membrane protein.; [Isoform 2]: Cell membrane; Single-pass type I membrane protein.; [Isoform 3]: Cytoplasm . Detected in the cytoplasm of spermatozoa, especially in the

equatorial and subacrosomal region of the sperm head.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

epitope-specific immunogen. chromatography using





Explore. Bioreagents.

**Clonality** Polyclonal

Concentration 1 mg/ml

Observed band 110kD

Human Gene ID 3815

Human Swiss-Prot Number P10721

Alternative Names KIT; SCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait


protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD

antigen CĎ117

Background This gene encodes the human homolog of the proto-oncogene c-kit. C-kit

was first identified as the cellular homolog of the feline sarcoma viral oncogene v-kit. This protein is a type 3 transmembrane receptor for MGF (mast cell growth factor, also known as stem cell factor). Mutations in this gene are associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous lukemia, and piebaldism. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jul 2008],




Western Blot analysis of various cells using Phospho-c-Kit (Y721) Polyclonal Antibody






Explore. Bioreagents.



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using c-Kit (Phospho-Tyr721) Antibody



Western blot analysis of lysates from HepG2 cells treated with EGF 200ng/ml 30', using c-Kit (Phospho-Tyr721) Antibody. The lane on the right is blocked with the phospho peptide.