

eIF3K rabbit pAb**Cat#: orb765120 (Manual)**

For research use only. Not intended for diagnostic use.

Product Name	eIF3K rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human EIF3K. AA range:61-110
Specificity	eIF3K Polyclonal Antibody detects endogenous levels of eIF3K protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Eukaryotic translation initiation factor 3 subunit K
Gene Name	EIF3K
Cellular localization	Nucleus . Cytoplasm .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal

Concentration 1 mg/ml

Observed band 30kD

Human Gene ID 27335


Human Swiss-Prot Number Q9UBQ5

Alternative Names EIF3K; EIF3S12; ARG134; HSPC029; MSTP001; PTD001; Eukaryotic translation initiation factor 3 subunit K; eIF3k; Eukaryotic translation initiation factor 3 subunit 12; Muscle-specific gene M9 protein; PLAC-24; eIF-3 p25; eIF-3 p28

Background The 700-kD eukaryotic translation initiation factor-3 (eIF3) is the largest eIF and contains at least 12 subunits, including EIF2S12. eIF3 plays an essential role in translation by binding directly to the 40S ribosomal subunit and promoting formation of the 40S preinitiation complex (Mayeur et al., 2003 [PubMed 14519125]).[supplied by OMIM, Mar 2008],



Western Blot analysis of various cells using eIF3K Polyclonal Antibody diluted at 1:1000



Western blot analysis of lysates from HeLa cells, using EIF3K Antibody. The lane on the right is blocked with the synthesized peptide.