

Histone deacetylase 9 rabbit pAb**Cat#: orb765397 (Manual)**


For research use only. Not intended for diagnostic use.

Product Name	Histone deacetylase 9 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human HDAC9. AA range:1017-1066
Specificity	Histone deacetylase 9 Polyclonal Antibody detects endogenous levels of Histone deacetylase 9 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Histone deacetylase 9
Gene Name	HDAC9
Cellular localization	Nucleus .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal


Concentration	1 mg/ml
Observed band	110kD
Human Gene ID	9734
Human Swiss-Prot Number	Q9UKV0-5,6,7
Alternative Names	Histone deacetylase 9; HD9; Histone deacetylase 7B; HD7; HD7b; Histone deacetylase-related protein; MEF2-interacting transcription repressor MITR; HDAC9; HDAC7; HDAC7B; HDRP; KIAA0744; MITR

Background


Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. Histone deacetylase 9 encoded by HDAC9 has sequence homology to members of the histone deacetylase family. HDAC9 is orthologous to the Xenopus and mouse MITR genes. The MITR protein lacks the histone deacetylase catalytic domain. It represses MEF2 activity through recruitment of multicomponent corepressor complexes that include CtBP and HDACs. This encoded protein may play a role in hematopoiesis. Multiple alternatively spliced transcripts have been described for this gene but the full-length nature of some of them has not been determined.




Western Blot analysis of various cells using Histone deacetylase 9 Polyclonal Antibody diluted at 1:1000



Western Blot analysis of HepG2 cells using Histone deacetylase 9 Polyclonal Antibody diluted at 1:1000



Immunofluorescence analysis of HepG2 cells, using HDAC9 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using HDAC9 Antibody. The picture on the right is blocked with the synthesized peptide.