

MDA5 rabbit pAb**Cat#: orb765637 (Manual)**


For research use only. Not intended for diagnostic use.

Product Name	MDA5 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human IFIH1. AA range:976-1025
Specificity	MDA5 Polyclonal Antibody detects endogenous levels of MDA5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Interferon-induced helicase C domain-containing protein 1
Gene Name	IFIH1
Cellular localization	Cytoplasm . Nucleus . Mitochondrion . Upon viral RNA stimulation and ISGylation, translocates from cytosol to mitochondrion. May be found in the nucleus, during apoptosis.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.


Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	120kD
Human Gene ID	64135
Human Swiss-Prot Number	Q9BYX4
Alternative Names	IFIH1; MDA5; RH116; Interferon-induced helicase C domain-containing protein 1; Clinically amyopathic dermatomyositis autoantigen 140 kDa; CADM-140 autoantigen; Helicase with 2 CARD domains; Helicard; Interferon-induced with helicase C domai

Background


DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein that is upregulated in response to treatment with beta-interferon and a protein kinase C-activating compound, mezerein. Irreversible reprogramming of melanomas can be achieved by treatment with both these agents; treatment with either agent alone only achieves reversible differentiation. Genetic variation in this gene is associated with diabetes mellitus insulin-depend



Western Blot analysis of various cells using MDA5 Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorb



Western blot analysis of lysates from Jurkat cells, using IFIH1 Antibody. The lane on the right is blocked with the synthesized peptide.