



## PIAS 1 rabbit pAb

Cat#: orb766078 (Manual)

For research use only. Not intended for diagnostic use.

Product Name PIAS 1 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA:

1/10000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human PIAS1. AA range:10-59

Specificity PIAS 1 Polyclonal Antibody detects endogenous levels of PIAS 1 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name E3 SUMO-protein ligase PIAS1

Gene Name PIAS1

Cellular localization Nucleus speckle . Nucleus, PML body . Interaction with CSRP2 may induce

a partial redistribution along the cytoskeleton.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal





Concentration 1 mg/ml

**Observed band** 72kD

**Human Gene ID** 8554

**Human Swiss-Prot Number** O75925


PIAS1; DDXBP1; E3 SUMO-protein ligase PIAS1; DEAD/H box-binding protein 1; Gu-binding protein; GBP; Protein inhibitor of activated STAT **Alternative Names** 

protein 1; RNA helicase II-binding protein


**Background** 

This gene encodes a member of the protein inhibitor of activated STAT (PIAS) family. PIAS proteins function as SUMO E3 ligases and play important roles in many cellular processes by mediating the sumoylation of target proteins. This protein plays a central role as a transcriptional

coregulator of numerous cellular pathways includign the STAT1 and nuclear factor kappaB pathways. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],



Western Blot analysis of various cells using PIAS 1 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).




Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using PIAS1 Antibody. The picture on the right is blocked with the synthesized peptide.





Explore. Bioreagents.



Western blot analysis of lysates from HUVEC cells, using PIAS1 Antibody. The lane on the right is blocked with the synthesized peptide.