



NFAT5 (phospho Ser1197) rabbit pAb

Cat#: orb767415 (Manual)

For research use only. Not intended for diagnostic use.

Product Name NFAT5 (phospho Ser1197) rabbit pAb

Host species Rabbit

Applications WB;IHC

Species Cross-Reactivity Human; Mouse

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300

Immunogen The antiserum was produced against synthesized peptide derived from

human NFAT5 around the phosphorylation site of Ser1197. AA range:1171-

1220

Specificity Phospho-NFAT5 (S1197) Polyclonal Antibody detects endogenous levels of

NFAT5 protein only when phosphorylated at \$1197.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Nuclear factor of activated T-cells 5

Gene Name NFAT5

Cellular localization Nucleus . Cytoplasm . Nuclear distribution increases under hypertonic

conditions.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Explore. Bioreagents.

Concentration 1 mg/ml

Observed band

Human Gene ID 10725


Human Swiss-Prot Number O94916

NFAT5; KIAA0827; TONEBP; Nuclear factor of activated T-cells 5; NFAT5; T-cell transcription factor NFAT5; Tonicity-responsive enhancer-binding protein; TonE-binding protein; TonEBP **Alternative Names**


Background The product of this gene is a member of the nuclear factors of activated T

cells family of transcription factors. Proteins belonging to this family play a central role in inducible gene transcription during the immune response. This protein regulates gene expression induced by osmotic stress in mammalian cells. Unlike monomeric members of this protein family, this protein exists as a homodimer and forms stable dimers with DNA elements. Multiple transcript variants encoding different isoforms have been found for this gene.

[provided by RefSeq, Jul 2008],




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using NFAT5 (Phospho-Ser1197) Antibody



Immunofluorescence analysis of COS7 cells, using NFAT5 (Phospho-Ser1197) Antibody. The picture on the right is blocked with the phospho peptide.







Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).