

Adenosine A2A-R rabbit pAb**Cat#: orb767652 (Manual)**


For research use only. Not intended for diagnostic use.

Product Name	Adenosine A2A-R rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human ADORA2A. AA range: 120-169
Specificity	Adenosine A2A-R Polyclonal Antibody detects endogenous levels of Adenosine A2A-R protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Adenosine receptor A2a
Gene Name	ADORA2A
Cellular localization	Cell membrane ; Multi-pass membrane protein . Colocalizes with GAS2L2 at neuronal processes. .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal


Concentration	1 mg/ml
Observed band	37kD
Human Gene ID	135
Human Swiss-Prot Number	P29274
Alternative Names	ADORA2A; ADORA2; Adenosine receptor A2a

Background


adenosine A2a receptor(ADORA2A) Homo sapiens This gene encodes a member of the guanine nucleotide-binding protein (G protein)-coupled receptor (GPCR) superfamily, which is subdivided into classes and subtypes. The receptors are seven-pass transmembrane proteins that respond to extracellular cues and activate intracellular signal transduction pathways. This protein, an adenosine receptor of A2A subtype, uses adenosine as the preferred endogenous agonist and preferentially interacts with the G(s) and G(olf) family of G proteins to increase intracellular cAMP levels. It plays an important role in many biological functions, such as cardiac rhythm and circulation, cerebral and renal blood flow, immune function, pain regulation, and sleep. It has been implicated in pathophysiological conditions such as inflammatory diseases and neurodegenerative disorders. Alternative splicing results in multiple transcript variants. A read-through transcript compos




Immunofluorescence analysis of COS7 cells, using ADORA2A Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using ADORA2A Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of ADORA2A Antibody. The lane on the right is blocked with the ADORA2A peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using ADORA2A Antibody. The lane on the right is blocked with the ADORA2A peptide.