



## NT5C3 rabbit pAb

**Cat#: orb769381 (Manual)** 

For research use only. Not intended for diagnostic use.

Product Name NT5C3 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human NT5C3. AA range:11-60

Specificity NT5C3 Polyclonal Antibody detects endogenous levels of NT5C3 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Cytosolic 5'-nucleotidase 3

Gene Name NT5C3

Cellular localization Cytoplasm .; [Isoform 2]: Endoplasmic reticulum.

**Purification** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Clonality** Polyclonal





Explore. Bioreagents.

1 mg/ml Concentration

**Observed band** 38kD


**Human Gene ID** 51251

**Human Swiss-Prot Number** Q9H0P0

NT5C3; P5N1; UMPH1; HSPC233; Cytosolic 5'-nucleotidase 3; Cytosolic 5'-nucleotidase III; cN-III; Pyrimidine 5'-nucleotidase 1; P5'N-1; P5N-1; PN-I; Uridine 5'-monophosphate hydrolase 1; p36 **Alternative Names** 

5'-nucleotidase, cytosolic IIIA(NT5C3A) Homo sapiens This gene encodes a member of the 5'-nucleotidase family of enzymes that catalyze the **Background** 

dephosphorylation of nucleoside 5'-monophosphates. The encoded protein is the type 1 isozyme of pyrimidine 5' nucleotidase and catalyzes the dephosphorylation of pyrimidine 5' monophosphates. Mutations in this gene are a cause of hemolytic anemia due to uridine 5-prime monophosphate hydrolase deficiency. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and pseudogenes of this gene are located on the long arm of chromosomes 3 and 4. [provided by RefSeq, Mar 2012],




Immunohistochemistry analysis of paraffin-embedded human brain tissue, using NT5C3 Antibody. The picture on the right is blocked with the synthesized peptide.






Explore. Bioreagents.



Western blot analysis of lysates from Jurkat and COLO205 cells, using NT5C3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from 293 cells using NT5C3 antibody.