



MRP-L39 rabbit pAb

Cat#: orb769517 (Manual)

For research use only. Not intended for diagnostic use.

Product Name MRP-L39 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA:

1/40000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human MRPL39. AA range:289-338

Specificity MRP-L39 Polyclonal Antibody detects endogenous levels of MRP-L39

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name 39S ribosomal protein L39 mitochondrial

Gene Name MRPL39

Cellular localization Mitochondrion .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





1 mg/ml Concentration

Observed band 39kD

Human Gene ID 54148

Human Swiss-Prot Number Q9NYK5

MRPL39; C21orf92; MRPL5; RPML5; MSTP003; PRED22; 39S ribosomal protein L39; mitochondrial; L39mt; MRP-L39; 39S ribosomal protein L5, **Alternative Names**


mitochondrial; L5mt; MRP-L5

Background Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes

and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 39S subunit protein. Two transcript variants encoding distinct isoforms have been described. A pseudogene

corresponding to this gene is found on chromosome 5q. [provided by


RefSeq, Jul 2008],




Western Blot analysis of 3T3 cells using MRP-L39 Polyclonal Antibody diluted at 1:1000







Immunohistochemistry analysis of paraffin-embedded human heart tissue, using MRPL39 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat and MCF-7 cells, using MRPL39 Antibody. The lane on the right is blocked with the synthesized peptide.