



PKD1 (phospho Ser910) rabbit pAb

Cat#: orb769660 (Manual)

For research use only. Not intended for diagnostic use.

Product Name PKD1 (phospho Ser910) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human PKD1/PKC mu around the phosphorylation site of Ser910. AA

range:863-912

Specificity Phospho-PKD1 (S910) Polyclonal Antibody detects endogenous levels of

PKD1 protein only when phosphorylated at S910.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Serine/threonine-protein kinase D1

Gene Name PRKD1

Cellular localization Cytoplasm . Cell membrane . Golgi apparatus, trans-Golgi network .

Translocation to the cell membrane is required for kinase activation.

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Concentration

Observed band 117kD

1 mg/ml

Human Gene ID 5587


Human Swiss-Prot Number Q15139

PRKD1; PKD; PKD1; PRKCM; Serine/threonine-protein kinase D1; Protein **Alternative Names**


kinase C mu type; Protein kinase D; nPKC-D1; nPKC-mu

Background PRKD1 is a serine/threonine kinase that regulates a variety of cellular

functions, including membrane receptor signaling, transport at the Golgi, protection from oxidative stress at the mitochondria, gene transcription, and regulation of cell shape, motility, and adhesion (summary by Eiseler et al., 2009 [PubMed 19329994]).[supplied by OMIM, Nov 2010],



Western Blot analysis of A431 cells using Phospho-PKD1 (S910) Polyclonal Antibody




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PKD1/PKC mu (Phospho-Ser910) Antibody





Explore. Bioreagents.



Western blot analysis of lysates from A431 cells, using PKD1/PKC mu (Phospho-Ser910) Antibody. The lane on the right is blocked with the phospho peptide.



Immunohistochemical analysis of paraffin-embedded human small intestinal carcinoma tissue. 1,primary Antibody was diluted at 1:200(4° overnight). 2, Sodium citrate pH 6.0 was used for antigen retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200