



PKR (phospho Thr446) rabbit pAb

Cat#: orb769711 (Manual)

For research use only. Not intended for diagnostic use.

Product Name PKR (phospho Thr446) rabbit pAb

Host species Rabbit

Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human PKR around the phosphorylation site of Thr446. AA range:413-462

Specificity Phospho-PKR (T446) Polyclonal Antibody detects endogenous levels of

PKR protein only when phosphorylated at T446.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Interferon-induced double-stranded RNA-activated protein kinase

Gene Name EIF2AK2

Cellular localization Cytoplasm . Nucleus . Cytoplasm, perinuclear region . Nuclear localization is

elevated in acute leukemia, myelodysplastic syndrome (MDS), melanoma, breast, colon, prostate and lung cancer patient samples or cell lines as well as

neurocytes from advanced C

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.





Explore. Bioreagents.

Clonality Polyclonal

Concentration 1 mg/ml

Observed band 62kD

Human Gene ID 5610

Human Swiss-Prot Number P19525

Alternative Names EIF2AK2; PKR; PRKR; Interferon-induced; double-stranded RNA-


activated protein kinase; Eukaryotic translation initiation factor 2-alpha kinase 2; eIF-2A protein kinase 2; Interferon-inducible RNA-dependent

protein kinase; P1/eIF-2A protein k

Background The protein encoded by this gene is a serine/threonine protein kinase that is

activated by autophosphorylation after binding to dsRNA. The activated form of the encoded protein can phosphorylate translation initiation factor EIF2S1, which in turn inhibits protein synthesis. This protein is also activated by manganese ions and heparin. Three transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq,

Oct 2011],




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PKR (Phospho-Thr446) Antibody





Explore. Bioreagents.



Western blot analysis of lysates from NIH/3T3 cells treated with IFN 2500U/ml 30', using PKR (Phospho-Thr446) Antibody. The lane on the right is blocked with the phospho peptide.