



Raf-B (phospho Thr599) rabbit pAb

Cat#: orb770156 (Manual)

For research use only. Not intended for diagnostic use.

Product Name Raf-B (phospho Thr599) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions WB 1:500-2000 ,Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000.

Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized peptide derived from

human B-RAF around the phosphorylation site of Thr599. AA range:571-

620

Specificity Phospho-Raf-B (T599) Polyclonal Antibody detects endogenous levels of

Raf-B protein only when phosphorylated at T599.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Serine/threonine-protein kinase B-raf

Gene Name BRAF

Cellular localization Nucleus . Cytoplasm . Cell membrane . Colocalizes with RGS14 and RAF1

in both the cytoplasm and membranes. .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Clonality Polyclonal





Explore. Bioreagents.

Concentration 1 mg/ml

Observed band

Human Gene ID 673

Human Swiss-Prot Number P15056


Alternative Names BRAF; BRAF1; RAFB1; Serine/threonine-protein kinase B-raf; Proto-

oncogene B-Raf; p94; v-Raf murine sarcoma viral oncogene homolog B1

Background This gene encodes a protein belonging to the raf/mil family of

serine/threonine protein kinases. This protein plays a role in regulating the MAP kinase/ERKs signaling pathway, which affects cell division, differentiation, and secretion. Mutations in this gene are associated with cardiofaciocutaneous syndrome, a disease characterized by heart defects, mental retardation and a distinctive facial appearance. Mutations in this gene have also been associated with various cancers, including non-Hodgkin lymphoma, colorectal cancer, malignant melanoma, thyroid carcinoma, non-small cell lung carcinoma, and adenocarcinoma of lung. A pseudogene, which is located on chromosome X, has been identified for this gene.


[provided by RefSeq, Jul 2008],



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using B-RAF (Phospho-Thr599) Antibody







Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using B-RAF (Phospho-Thr599) Antibody. The picture on the right is blocked with the phospho peptide.