

**TNF-IP 8L2 rabbit pAb****Cat#: orb770461 (Manual)**


For research use only. Not intended for diagnostic use.

<b>Product Name</b>	TNF-IP 8L2 rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;ELISA
<b>Species Cross-Reactivity</b>	Human;Mouse;Rat
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human TNFAIP8L2. AA range:11-60
<b>Specificity</b>	TNF-IP 8L2 Polyclonal Antibody detects endogenous levels of TNF-IP 8L2 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	Tumor necrosis factor alpha-induced protein 8-like protein 2
<b>Gene Name</b>	TNFAIP8L2
<b>Cellular localization</b>	cytoplasm,
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal

<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	20kD
<b>Human Gene ID</b>	79626
<b>Human Swiss-Prot Number</b>	Q6P589
<b>Alternative Names</b>	TNFAIP8L2; Tumor necrosis factor alpha-induced protein 8-like protein 2; TIPE2; TNF alpha-induced protein 8-like protein 2; TNFAIP8-like protein 2; Inflammation factor protein 20

## Background

domain:The central region was initially thought to constitute a DED (death effector) domain. However, 3D-structure data reveal a previously uncharacterized fold that is different from the predicted fold of a DED (death effector) domain. It consists of a large, hydrophobic central cavity that is poised for cofactor binding.,function:Acts as a negative regulator of innate and adaptive immunity by maintaining immune homeostasis. Negative regulator of Toll-like receptor and T-cell receptor function. Prevents hyperresponsiveness of the immune system and maintains immune homeostasis. Inhibits jun/ap1 and NF-kappa-B activation. Promotes Fas-induced apoptosis.,similarity:Belongs to the TNFAIP8 family. TNFAIP8L2 subfamily.,subunit:May interact with CASP8; however, such result is unclear since PubMed:19079267 could not reproduce the interaction with CASP8.,



Western blot analysis of the lysates from HeLa cells using TNFAIP8L2 antibody.