

IKK γ (phospho Ser31) rabbit pAb**Cat#: orb770677 (Manual)**

For research use only. Not intended for diagnostic use.


Product Name	IKK γ (phospho Ser31) rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human IKK-gamma around the phosphorylation site of Ser31. AA range: 16-65
Specificity	Phospho-IKK γ (S31) Polyclonal Antibody detects endogenous levels of IKK γ protein only when phosphorylated at S31.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	NF-kappa-B essential modulator
Gene Name	IKBKG
Cellular localization	Cytoplasm . Nucleus . Sumoylated NEMO accumulates in the nucleus in response to genotoxic stress. .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal

Explore. Bioreagents.


Concentration	1 mg/ml
Observed band	48kD
Human Gene ID	8517
Human Swiss-Prot Number	Q9Y6K9
Alternative Names	IKBK γ ; FIP3; NEMO; NF-kappa-B essential modulator; NEMO; FIP-3; I κ B kinase-associated protein 1; IKKAP1; Inhibitor of nuclear factor kappa-B kinase subunit gamma; I-kappa-B kinase subunit gamma; IKK-gamma; IKK γ ; I κ B kinase subunit gamma; NF

Background


This gene encodes the regulatory subunit of the inhibitor of kappaB kinase (IKK) complex, which activates NF-kappaB resulting in activation of genes involved in inflammation, immunity, cell survival, and other pathways. Mutations in this gene result in incontinentia pigmenti, hypohidrotic ectodermal dysplasia, and several other types of immunodeficiencies. A pseudogene highly similar to this locus is located in an adjacent region of the X chromosome. [provided by RefSeq, Mar 2016],



Western Blot analysis of 293 cells using Phospho-IKK γ (S31) Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using IKK-gamma (Phospho-Ser31) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells treated with TNF- α 20ng/ml 5', using IKK-gamma (Phospho-Ser31) Antibody. The lane on the right is blocked with the phospho peptide.