

MDC1 (phospho Ser513) rabbit pAb**Cat#: orb770956 (Manual)**


For research use only. Not intended for diagnostic use.

Product Name	MDC1 (phospho Ser513) rabbit pAb
Host species	Rabbit
Applications	IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human MDC1 around the phosphorylation site of Ser513. AA range:479-528
Specificity	Phospho-MDC1 (S513) Polyclonal Antibody detects endogenous levels of MDC1 protein only when phosphorylated at S513.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Mediator of DNA damage checkpoint protein 1
Gene Name	MDC1
Cellular localization	Nucleus . Chromosome . Associated with chromatin. Relocalizes to discrete nuclear foci following DNA damage, this requires 'Ser-139' phosphorylation of H2AX. Colocalizes with APTX at sites of DNA double-strand breaks.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.


Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	
Human Gene ID	9656
Human Swiss-Prot Number	Q14676
Alternative Names	MDC1; KIAA0170; NFBD1; Mediator of DNA damage checkpoint protein 1; Nuclear factor with BRCT domains 1

Background

The protein encoded by this gene contains an N-terminal forkhead domain, two BRCA1 C-terminal (BRCT) motifs and a central domain with 13 repetitions of an approximately 41-amino acid sequence. The encoded protein is required to activate the intra-S phase and G2/M phase cell cycle checkpoints in response to DNA damage. This nuclear protein interacts with phosphorylated histone H2AX near sites of DNA double-strand breaks through its BRCT motifs, and facilitates recruitment of the ATM kinase and meiotic recombination 11 protein complex to DNA damage foci. [provided by RefSeq, Jul 2008],



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using MDC1 (Phospho-Ser513) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using MDC1 (Phospho-Ser513) Antibody. The picture on the right is blocked with the phospho peptide.