


**TRIF rabbit pAb****Cat#: orb771376 (Manual)**


For research use only. Not intended for diagnostic use.

<b>Product Name</b>	TRIF rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Species Cross-Reactivity</b>	Human;Rat;Mouse;
<b>Recommended dilutions</b>	WB 1:500-2000,IHC-p 1:50-200 ELISA 1:10000-20000
<b>Immunogen</b>	Synthesized peptide derived from TRIF at AA range: 663-712
<b>Specificity</b>	TRIF Polyclonal Antibody detects endogenous levels of TRIF
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	TRIF
<b>Gene Name</b>	TICAM1
<b>Cellular localization</b>	Cytoplasmic vesicle, autophagosome . Cytoplasm, cytosol . Mitochondrion . Colocalizes with UBQLN1 in the autophagosome (PubMed:21695056). Colocalizes in the cytosol with DDX1, DDX21 and DHX36. Colocalizes in the mitochondria with DDX1 and poly(I:C) RNA ligand. The multi-helicase-TICAM1 complex may translocate to the mitochondria upon poly(I:C) RNA ligand stimulation (By similarity). .


<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	80kD
<b>Human Gene ID</b>	148022
<b>Human Swiss-Prot Number</b>	Q8IUC6
<b>Alternative Names</b>	TIR domain-containing adapter molecule 1 (TICAM-1;Proline-rich, vinculin and TIR domain-containing protein B;Putative NF-kappa-B-activating protein 502H;Toll-interleukin-1 receptor domain-containing adapter protein inducing interferon beta;TIR domain-cont
<b>Background</b>	This gene encodes an adaptor protein containing a Toll/interleukin-1 receptor (TIR) homology domain, which is an intracellular signaling domain that mediates protein-protein interactions between the Toll-like receptors (TLRs) and signal-transduction components. This protein is involved in native immunity against invading pathogens. It specifically interacts with toll-like receptor 3, but not with other TLRs, and this association mediates dsRNA induction of interferon-beta through activation of nuclear factor kappa-B, during an antiviral immune response. [provided by RefSeq, Jan 2012],




**Western Blot analysis of HEPG2 cells using TRIF Polyclonal Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000**



**Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200**



**Immunohistochemical analysis of paraffin-embedded human-lung-cancer, antibody was diluted at 1:200**



**Immunohistochemical analysis of paraffin-embedded human-spleen, antibody was diluted at 1:200**