



CTGF rabbit pAb

Cat#: orb771624 (Manual)

For research use only. Not intended for diagnostic use.

Product Name CTGF rabbit pAb

Host species Rabbit

Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions IHC-p 1:50-200, ELISA 1:10000-20000

Immunogen Synthetic peptide from human protein at AA range: 141-190

The antibody detects endogenous CTGF **Specificity**

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium

azide..

Store at -20°C. Avoid repeated freeze-thaw cycles. **Storage**

Protein Name Connective tissue growth factor (CCN family member 2) (Hypertrophic

chondrocyte-specific protein 24) (Insulin-like growth factor-binding protein 8) (IBP-8) (IGF-binding protein 8) (IGFBP-8)

CTGF CCN2 HCS24 IGFBP8 Gene Name

Cellular localization Secreted, extracellular space, extracellular matrix . Secreted .

Purification The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.





Explore. Bioreagents.

Clonality Polyclonal

Concentration 1 mg/ml

Observed band

1490 **Human Gene ID**


Human Swiss-Prot Number P29279

Connective tissue growth factor (CCN family member 2;Hypertrophic chondrocyte-specific protein 24;Insulin-like growth factor-binding protein 8;IBP-8;IGF-binding protein 8;IGFBP-8) **Alternative Names**


Background

The protein encoded by this gene is a mitogen that is secreted by vascular endothelial cells. The encoded protein plays a role in chondrocyte proliferation and differentiation, cell adhesion in many cell types, and is related to platelet-derived growth factor. Certain polymorphisms in this gene have been linked with a higher incidence of systemic sclerosis. [provided by Profice New 2000]

RefSeq, Nov 2009],




Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:200







Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:200