

**NDUB8 rabbit pAb****Cat#: orb772751 (Manual)**

For research use only. Not intended for diagnostic use.

<b>Product Name</b>	NDUB8 rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;ELISA
<b>Species Cross-Reactivity</b>	Human;Mouse
<b>Recommended dilutions</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 120-200
<b>Specificity</b>	NDUB8 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide..
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 8, mitochondrial (Complex I-ASHI) (CI-ASHI) (NADH-ubiquinone oxidoreductase ASHI subunit)
<b>Gene Name</b>	NDUFB8
<b>Cellular localization</b>	Mitochondrion inner membrane ; Single-pass membrane protein ; Matrix side .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

**Clonality** Polyclonal

**Concentration** 1 mg/ml

**Observed band** 20kD


**Human Gene ID** 4714

**Human Swiss-Prot Number** O95169

**Alternative Names**

## Background

function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone. similarity: Belongs to the complex I NDUF8 subunit family. subunit: Complex I is composed of 45 different subunits.



**Western blot analysis of lysates from PC12 cells, primary antibody was diluted at 1:1000, 4° over night**