

Product Datasheet

Anti-Hu CD77 PE (orb1088485)

Description	CD77 (globotriaosylceramide Gb3), also known as the Pk blood group antigen, BLA (Burkitt's lymphoma associated antigen), or CTH (ceramide trihexoside) is a neutral glycosphingolipid composed of three carbohydrate molecules linked to a lipid moiety in the cell membrane (Gal-alpha1-4Gal-beta1-4Glc-beta1-Cer). It is expressed on germinal center B cells, Burkitt's lymphoma cells, it can be induced on extrafollicular B cells and it is also found on endothelia and epithelia. CD77 may be involved in elimination of germinal center B cells that fail to produce high affinity antibodies, and serves also as receptor for shiga toxin and verotoxin.
Reactivity	Human
Conjugation	PE
Tested Applications	FC
Immunogen	Daudi cell line (Burkitt's lymphoma)
Target	CD77
Preservatives	Stabilizing Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Storage	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.
Note	For research use only
Application notes	Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.
Isotype	Lewis Rat IgM
Clonality	Monoclonal
Clone Number	38.13

Biorbyt Ltd.


7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Purity Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Expiration Date 12 months from date of receipt.



Separation of RAJI cells stained using anti-human CD77 (38.13) PE antibody (10 μ l reagent per million cells in 100 μ l of cell suspension, red-filled) from RAJI cells stained using mouse IgG1 isotype control (MOPC-21) APC antibody (concentration in sample 5 μ g/ml, same as CD77 PE concentration, black-dashed) in flow cytometry analysis (surface staining) of RAJI cell suspension.

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)