


Product Datasheet

CD77 Antibody (FITC) (orb1088486)

Description	CD77 (globotriaosylceramide Gb3), also known as the Pk blood group antigen, BLA (Burkitt's lymphoma...
Reactivity	Human
Conjugation	FITC
Tested Applications	FC
Immunogen	Daudi cell line (Burkitt's lymphoma)
Target	CD77
Preservatives	Stabilizing Tris buffered saline (TBS), pH 8.0, 15 mM sodium azide
Storage	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.
Note	For research use only
Application notes	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.
Isotype	Lewis Rat IgM
Clonality	Monoclonal
Purity	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.
Dilution Range	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.
Expiration Date	12 months from date of receipt.



Separation of RAJI cells (red-filled) fr...