



## **Product Datasheet**

## **UBE2C Peptide (orb1235512)**

**Description** UBE2C Peptide

**Tested** Blocking

**Applications** 

Target UBE2C

**Preservatives** UBE2C peptide is supplied as 100 μg of dried

peptide. When the peptide is reconstituted in 200  $\mu L$  water, the concentration will be 0.5

mg/mL.

Form/Appearance Lyophilized

**Storage** Store at -20°C. When reconstituted minimize

freezing and thawing.

**Note** For research use only

**Application notes** UBE2C Peptide is used for blocking the activity

of UBE2C antibody (catalog number 42-945). The UBE2C antibody ships at 0.5 mg/mL and the UBE2C Peptide ships as a 100  $\mu$ g pellet. When peptide is reconstituted in 200  $\mu$ L water, the concentration will be 0.5 mg/mL, identical to that of the antibody. The best starting ratio would be 1:1 (which means molar excess of peptide relative to antibody when identical volumes are mixed). Mix equal volume of peptide and antibody at the required dilution and leave at ambient temperature for 20 minutes. Then incubate and develop in parallel two blots: one with the antibody and one with the antibody which has been pre-absorbed with

the provided peptide.

Uniprot ID 000762

NCBI NP\_861517.1, NP\_008950.1, NP\_861515.1

**Expiration Date** 6 months from date of receipt.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom