



## **Product Datasheet**

## **USP16 Peptide (orb1236202)**

**Description** USP16 Peptide

**Tested** Blocking

**Applications** 

Target USP16

**Preservatives** USP16 peptide is supplied as 100 μg of dried

peptide. When the peptide is reconstituted in 200  $\mu L$  water, the concentration will be 0.5

mg/mL.

Form/Appearance Lyophilized

**Storage** Store at -20°C. When reconstituted minimize

freezing and thawing.

**Note** For research use only

**Application notes** USP16 Peptide is used for blocking the activity

of USP16 antibody (catalog number 42-222). The USP16 antibody ships at 0.5 mg/mL and the USP16 Peptide ships as a 100 ug pellet. When peptide is reconstituted in 200 uL water, the concentration will be 0.5 mg/mL, identical to that of the antibody. The best starting ratio would be 1:1 (which means molar excess of peptide relative to antibody when identical volumes are mixed). Mix equal volume of peptide and antibody at the required dilution and leave at ambient temperature for 20 minutes. Then incubate and develop in parallel two blots: one with the antibody and one with the antibody which has been pre-absorbed with

the provided peptide.

Uniprot ID Q9Y5T5

NCBI NP\_001001992.1, NP\_001027582.1

**Expiration Date** 6 months from date of receipt.