

## **Product Datasheet**

Goat Anti-Rabbit IgG (H&L) Antibody (FITC) (orb1236937)



## www.biorbyt.com

Descriptionnts. Goat Anti-Rabbit IgG (H&L) Antibody

Species/Host Goat

**Reactivity** Rabbit

Conjugation FITC

**Tested** FC, IF

**Applications** 

Immunogen Rabbit IgG - whole molecule

**Preservatives** 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, 0.01% (w/v) Sodium Azide; pH 7.2; 10 mg/mL Bovine

Serum Albumin

Form/Appearance Lyophilized

**Storage** Fluorescein Conjugated Affinity

Purified anti-Rabbit IgG can be stored vial at 4 °C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20 °C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4 °C as an undiluted liquid. Dilute only

prior to immediate use.

**Note** For research use only

**Application notes** Fluorescein Conjugated Affinity

Purified anti-Rabbit IgG is suitable for

immunomicroscopy and flow

cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Recommended dilutions are as follows, Flow Cytometry 1:2,000 - 1:10,000, IF Microscopy 1:500 - 1:2,500. (Optimal dilution should be determined by

user).

**Isotype** IgG

**Dilution Range** Fluorescein Conjugated Affinity

Purified anti-Rabbit IgG is suitable for

immunomicroscopy and flow

cytometry or FACS analysis as well as other antibody based fluorescent

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280

Biorbyt LLC.

68 TW Alexander Drive<br>Research Triangle Park<br>Durham, North Carolina<br/>br>27709. United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558