

---

## Product Datasheet

### Goat Anti-Rabbit IgG (H&L) Antibody Pre-Adsorbed (orb1236940)

**Description**

Goat Anti-Rabbit IgG (H&amp;L) Antibody

<b>Species/Host</b>	Goat
<b>Reactivity</b>	Rabbit
<b>Conjugation</b>	Unconjugated
<b>Tested Applications</b>	ELISA, IHC, WB
<b>Immunogen</b>	Rabbit IgG - whole molecule
<b>Preservatives</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, 0.01% (w/v) Sodium Azide; pH 7.2
<b>Form/Appearance</b>	Liquid
<b>Concentration</b>	2.0 mg/mL
<b>Storage</b>	Unconjugated Affinity Purified anti-Rabbit IgG can be stored vial at 4 °C prior to restoration. For extended storage aliquot contents and freeze at -20 °C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4 °C as an undiluted liquid. Dilute only prior to immediate use.
<b>Note</b>	For research use only
<b>Application notes</b>	Unconjugated Affinity Purified anti-Rabbit IgG is suitable for immunoblotting (western or dot blot), ELISA and immunohistochemistry. Recommended dilutions are as follows, ELISA 1:35,000 - 1:325,000, Western Blot 1:3,000 - 1:35,000, Immunohistochemistry 1:1,000 - 1:5,000. (Optimal dilution should be determined by user).
<b>Isotype</b>	IgG
<b>Dilution Range</b>	Unconjugated Affinity Purified anti-Rabbit IgG is suitable for immunoblotting (western or dot blot), ELISA and immunohistochemistry. Recommended dilutions are as follows, ELISA 1:35,000 - 1:325,000, Western Blot 1:3,000 - 1:35,000, Immunohistochemistry 1:1,000 - 1:5,000. (Optimal dilution should be determined by user).