

---

## Product Datasheet

### Pkdcc Antibody (orb1238577)

**Description**

Pkdcc Antibody

**Species/Host**

Rabbit

**Reactivity**

Human, Mouse, Rat

**Conjugation**

Unconjugated

**Tested**

ELISA, IF, IHC-P, WB

**Applications**
**Immunogen**

VLK antibody was raised against a 15 amino acid synthetic peptide from near the center of human VLK. The immunogen is located within amino acids 320 - 370 of VLK.

**Target**

Pkdcc

**Preservatives**

VLK Antibody is supplied in PBS containing 0.02% sodium azide.

**Form/Appearance**

Liquid

**Concentration**

1 mg/mL

**Storage**

VLK antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Note**

For research use only

**Application notes**

VLK antibody can be used for detection of VLK by Western blot at 0.5 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 5 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in rat samples and Immunofluorescence in rat samples. All other applications and species not yet tested.

**Isotype**

IgG

**Clonality**


Polyclonal

**MW**


Predicted: 32, 54 kDa Observed: 31, 55 kDa

**Uniprot ID**
[Q5RJI4](#)
**NCBI**
[NP\\_612379](#)
**Dilution Range**


VLK antibody can be used for detection of VLK by Western blot at 0.5 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 5 µg/mL. Antibody validated: Western Blot in human



Western blot analysis of VLK in human lu...



Immunohistochemistry of VLK in rat lung ...



Immunofluorescence of VLK in rat lung ti...