
Product Datasheet

TTR Antibody (orb1238610)

Description

TTR Antibody

Species/Host

Gallus

Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Tested Applications

ELISA, IF, IHC-P, WB

Immunogen

Transthyretin antibody was raised against a 17 amino acid synthetic peptide near the center of human Transthyretin. The immunogen is located within amino acids 40 - 90 of Transthyretin.

Target

TTR

Preservatives

Transthyretin Antibody is supplied in PBS containing 0.02% sodium azide.

Form/Appearance

Liquid

Concentration

1 mg/mL

Storage

Transthyretin antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Note

For research use only

Application notes

Transthyretin antibody can be used for detection of Transthyretin by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Isotype

IgY

Clonality


Polyclonal

MW


Predicted: 15 kDa Observed: 16 kDa

Uniprot ID
[P02766](#)
NCBI
[P02766](#)
Dilution Range


Transthyretin antibody can be used for detection of Transthyretin by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in



Western blot analysis of 125 ng of Trans...



Immunohistochemistry of Transthyretin in...



Immunofluorescence of Transthyretin in h...