
Product Datasheet

UVRAG Antibody (orb1238622)

Description

UVRAG Antibody

Species/Host

Rabbit

Reactivity

Human, Mouse

Conjugation

Unconjugated

Tested Applications

ELISA, IF, IHC-P, WB

Immunogen

UVRAG antibody was raised against an 18 amino acid peptide near the carboxy terminus of human UVRAG. The immunogen is located within amino acids 490 - 540 of UVRAG.

Target

UVRAG

Preservatives

UVRAG antibody is supplied in PBS containing 0.02% sodium azide.

Form/Appearance

Liquid

Concentration

1 mg/mL

Storage

UVRAG antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Note

For research use only

Application notes

UVRAG antibody can be used for detection of UVRAG by Western blot at 1 - 2 µg/ml. Antibody can also be used for Immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in mouse samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Isotype

IgG

Clonality


Polyclonal

MW


Predicted: 77 kDa Observed: 75 kDa

Uniprot ID
[Q9P2Y5](#)
NCBI
[NP_003360](#)
Dilution Range


UVRAG antibody can be used for detection of UVRAG by Western blot at 1 - 2 µg/ml. Antibody can also be used for Immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in mouse samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.



Western blot analysis of UVRAG in mouse ...



Immunohistochemistry of UVRAG in human I...



Immunofluorescence of UVRAG in human lun...