
Product Datasheet

TXNIP Antibody (orb1238645)

Description

TXNIP Antibody

Species/Host

Rabbit

Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Tested Applications

ELISA, IF, IHC-P, WB

Immunogen

TXNIP antibody was raised against a 16 amino acid peptide near the center of human TXNIP. The immunogen is located within amino acids 110 - 160 of TXNIP.

Target

TXNIP

Preservatives

TXNIP antibody is supplied in PBS containing 0.02% sodium azide.

Form/Appearance

Liquid

Concentration

1 mg/mL

Storage

TXNIP antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Note

For research use only

Application notes

TXNIP body can be used for detection of TXNIP by Western blot at 0.5 - 1 µg/ml. Antibody can also be used for Immunohistochemistry at 2 µg/mL. For Immunoflorescence start at 5 µg/mL. Antibody validated: Western Blot in mouse samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Isotype

IgG

Clonality


Polyclonal

MW


Predicted: 43 kDa Observed: 39 kDa

Uniprot ID
[Q9H3M7](#)
NCBI
[NP_006463](#)
Dilution Range


TXNIP body can be used for detection of TXNIP by Western blot at 0.5 - 1 µg/ml. Antibody can also be used for Immunohistochemistry at 2 µg/mL. For Immunoflorescence start at 5 µg/mL. Antibody validated: Western Blot in mouse samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.



Western blot analysis of TXNIP in C2C12 ...



Immunohistochemistry of TXNIP in human k...



Immunofluorescence of TXNIP in human kid...