

## **Product Datasheet**

TRIM71 Antibody (orb1238674)



## www.biorbyt.com


55-

**▼TRIM71** 72-

Western blot analysis of TRIM71 in human...



Immunohistochemistry of TRIM71 in human ...



**Immunofluorescence** of TRIM71 in Human Br...

Descriptionnts. TRIM71 Antibody

Species/Host Rabbit

Reactivity Human, Mouse, Rat

Conjugation Unconjugated

Tested ELISA, IF, IHC-P, WB

**Applications** 

**Immunogen** TRIM71 antibody was raised against a 15 amino acid

synthetic peptide near the carboxy terminus of human TRIM71. The immunogen is located within

amino acids 680 - 730 of TRIM71.

TRIM71 **Target** 

**Preservatives** TRIM71 Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/mL

**Storage** TRIM71 antibody can be stored at 4°C for three

months and -20°C, stable for up to one year. As with

all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not

be exposed to prolonged high temperatures.

Note For research use only

**Application notes** TRIM71 antibody can be used for detection of

> TRIM71 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5

μg/mL. For immunofluorescence start at 20 μg/mL.Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Isotype IgG

Clonality Polyclonal

**Uniprot ID Q2Q1W2** 

NCBI NP 001034200

**Dilution Range** TRIM71 antibody can be used for detection of

> TRIM71 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5

μg/mL. For immunofluorescence start at 20 μg/mL.Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples

Carolina < br > 27709. United States

68 TW Alexander Drive<br/>br>Research Triangle Park<br/>br>Durham, North