
Product Datasheet

TLR1 Antibody (orb1238764)

Description

TLR1 Antibody

Species/Host

Rabbit

Reactivity

Human, Mouse

Conjugation

Unconjugated

Tested Applications

ELISA, IF, IHC-P, WB

Immunogen

TLR1 antibody was raised against a peptide corresponding to 16 amino acids near the amino terminus of human TLR1. The immunogen is located within amino acids 160 - 210 of TLR1.

Target

TLR1

Preservatives

TLR1 Antibody is supplied in PBS containing 0.02% sodium azide.

Form/Appearance

Liquid

Concentration

1 mg/mL

Storage

TLR1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Note

For research use only

Application notes

TLR1 antibody can be used for detection of TLR1 by Western blot at 1 to 4 µg/mL. Antibody can also be used for immunohistochemistry starting at 10 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in mouse samples; Immunohistochemistry in mouse samples and Immunofluorescence in mouse samples. All other applications and species not yet tested.

Isotype

IgG

Clonality


Polyclonal

MW


Predicted: 86 kDa Observed: 90 kDa

Uniprot ID
[Q15399](#)
NCBI
[CAG38593](#)
Dilution Range


TLR1 antibody can be used for detection of TLR1 by Western blot at 1 to 4 µg/mL. Antibody can also be used for immunohistochemistry starting at 10 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in mouse samples;



Western blot analysis of TLR1 in mouse s...



Immunohistochemistry of TLR1 in mouse sp...



Immunofluorescence of TLR1 in Mouse Sple...