


Product Datasheet

TICAM2 Antibody (orb1238769)


Description	TICAM2 Antibody
Species/Host	Rabbit
Reactivity	Human, Mouse, Rat
Conjugation	Unconjugated
Tested Applications	ELISA, IF, IHC-P, WB
Immunogen	TIRP antibody was raised against a peptide corresponding to amino acids near the C-terminus of human TIRP. The immunogen is located within the last 50 amino acids of TIRP.
Target	TICAM2
Preservatives	TIRP Antibody is supplied in PBS containing 0.02% sodium azide.
Form/Appearance	Liquid
Concentration	1 mg/mL
Storage	TIRP antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Note	For research use only
Application notes	TIRP antibody can be used for detection of TIRP by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human, mouse, and rat samples; Immunohistochemistry in mouse samples and Immunofluorescence in mouse samples. All other applications and species not yet tested.
Isotype	IgG
Clonality	Polyclonal
Uniprot ID	Q86XR7
NCBI	AAP81748
Dilution Range	TIRP antibody can be used for detection of TIRP by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human, mouse, and rat samples; Immunohistochemistry in mouse samples



Western blot analysis of TIRP in human (...)



Immunohistochemical staining of mouse kidney



Immunofluorescence of TIRP in Mouse Kidney