

Product Datasheet


Pmaip1 Antibody (orb1238793)




Applications

www.biorbyt.com


A B C



Western blot analysis of Noxa in human s...



Immunohistochemistry of NOXA in human st...



Immunofluorescence of Noxa in Human Stom...

Description Pmaip1 Antibody

Species/Host Rabbit

Reactivity Human, Mouse, Rat

Conjugation Unconjugated

Tested ELISA, IF, IHC-P, WB

Immunogen Noxa antibody was raised against a synthetic peptide corresponding to 17 amino acids at the

amino terminus of mouse Noxa. The immunogen is located within the first 50 amino acids of Noxa.

Target Pmaip1

Preservatives Noxa Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/mL

Storage Noxa antibody can be stored at 4°C for three months

and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Note For research use only

Application notes Noxa antibody can be used for detection of Noxa by

Western blot at at 0.5 - 2 μ g/mL. Antibody can also be used for immunohistochemistry starting at 1 μ g/mL. For immunofluorescence start at 10 μ g/mL.Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Isotype IgG

Clonality Polyclonal

Uniprot ID Q9JM54

NCBI NP 067426

Dilution Range Noxa antibody can be used for detection of Noxa by

Western blot at at 0.5 - 2 μ g/mL. Antibody can also be used for immunohistochemistry starting at 1 μ g/mL. For immunofluorescence start at 10 μ g/mL.Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples

Carolina

27709. United States

68 TW Alexander Drive
br>Research Triangle Park
br>Durham, North