


Product Datasheet

C12orf5 Antibody (orb1238797)


Description	C12orf5 Antibody
Species/Host	Rabbit
Reactivity	Human, Mouse
Conjugation	Unconjugated
Tested Applications	ELISA, IF, IHC-P, WB
Immunogen	TIGAR antibody was raised against a 19 amino acid synthetic peptide from near the center of human TIGAR. The immunogen is located within amino acids 100 - 150 of TIGAR.
Target	C12orf5
Preservatives	TIGAR Antibody is supplied in PBS containing 0.02% sodium azide.
Form/Appearance	Liquid
Concentration	1 mg/mL
Storage	TIGAR antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Note	For research use only
Application notes	TIGAR antibody can be used for detection of TIGAR by Western blot at 0.5 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.
Isotype	IgG
Clonality	Polyclonal
Uniprot ID	Q9NQ88
NCBI	NP_065108
Dilution Range	TIGAR antibody can be used for detection of TIGAR by Western blot at 0.5 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples



Western blot analysis of TIGAR in MCF7 c...



Immunohistochemistry of TIGAR in human b...



Immunofluorescence of TIGAR in Human Bra...