


Product Datasheet

CFLAR Antibody (orb1238926)


Description	CFLAR Antibody
Species/Host	Rabbit
Conjugation	Unconjugated
Tested Applications	ELISA, ICC, IF, WB
Immunogen	FLIP antibody was raised against a peptide corresponding to amino acids near the C-terminus of human FLIPa/FLIPI form. The immunogen is located within the last 50 amino acids of FLIP.
Target	CFLAR
Preservatives	FLIP Antibody is supplied in PBS containing 0.02% sodium azide.
Form/Appearance	Liquid
Concentration	1 mg/ml
Storage	FLIP antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Note	For research use only
Application notes	Flip antibody can be used for detection of FLIP by Western blot at 1 µg/mL. A 55 kDa band can be detected. Antibody can also be used for immunocytochemistry starting at 10 µg/mL. For immunofluorescence start at 10 µg/mL. Antibody validated: Western Blot in human samples; Immunocytochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.
Isotype	IgG
Clonality	Polyclonal
MW	55 kDa
Uniprot ID	O15519
NCBI	AAC51622
Dilution Range	Flip antibody can be used for detection of FLIP by Western blot at 1 µg/mL. A 55 kDa band can be detected. Antibody can also be used for immunocytochemistry starting at 10 µg/mL. For immunofluorescence start at 10 µg/mL. Antibody



Western blot analysis of FLIP in HeLa (H...)



Immunocytochemistry of FLIP in Jurkat cel...



Immunofluorescence of FLIP in Jurkat cel...