

---

## Product Datasheet

### NAIP Antibody (orb1239408)

**Description**

NAIP Antibody

**Species/Host**

Rabbit

**Reactivity**

Human

**Conjugation**

Unconjugated

**Tested**

ELISA, ICC, IF, WB

**Applications**
**Immunogen**

NAIP antibody was raised against a synthetic peptide corresponding to 13 amino acids at the C-terminus of human NAIP. The immunogen is located within amino acids 1200 - 1250 of NAIP.

**Target**

NAIP

**Preservatives**

NAIP Antibody is supplied in PBS containing 0.02% sodium azide.

**Form/Appearance**

Liquid

**Concentration**

1 mg/mL

**Storage**

NAIP antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Note**

For research use only

**Application notes**

NAIP antibody can be used for the detection of NAIP by Western blot at 0.5 to 1 µg/mL. Antibody can also be used for immunocytochemistry starting at 10 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples; Immunocytochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

**Isotype**

IgG

**Clonality**


Polyclonal

**MW**


Predicted: 137, 154 kDa Observed: 155 kDa

**Uniprot ID**
[Q13075](#)
**NCBI**
[AAC52047](#)
**Dilution Range**


NAIP antibody can be used for the detection of NAIP by Western blot at 0.5 to 1 µg/mL. Antibody can also be used for immunocytochemistry starting at 10 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human



Western blot analysis of NAIP in human t...



Immunocytochemistry of NAIP in A549 cell...



Immunofluorescence of NAIP in A549 cells...