



Product Datasheet

IRAK3 Antibody (orb1239522)



Description IRAK3 Antibody

Species/Host Rabbit

Reactivity Human, Mouse, Rat

Conjugation Unconjugated

Tested ELISA, IF, IHC-P, WB

Applications

Immunogen IRAK-M antibody was raised against a peptide

corresponding to 16 amino acids near the carboxy terminus of human IRAK-M.The immunogen is located within the last 50 amino acids of IRAK-M.

Target IRAK3

Preservatives IRAK-M Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/mL

Storage IRAK-M antibody can be stored at 4°C for three

months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated

freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Note For research use only

Application notes IRAK-M antibody can be used for detection of IRAK-M

by Western blot at 1 μ g/mL. IRAK-M antibody can also detect IRAK-M by immunohistochemistry at 2 μ g/mL. For immunofluorescence start at 10 μ g/mL.Antibody validated: Western Blot in mouse and rat samples; Immunohistochemistry in human and rat samples and Immunofluorescence in mouse and rat samples. All other applications and species

not yet tested.

Isotype IgG

Clonality Polyclonal

MW Predicted: 66 kDa Observed: 70 kDa

Uniprot ID Q9Y616

NCBI NP_009130

Dilution Range IRAK-M antibody can be used for detection of IRAK-M


by Western blot at 1 $\mu g/mL$. IRAK-M antibody can also detect IRAK-M by immunohistochemistry at 2

ualml For immunofluoroscopco start at 10


www.biorbyt.com

M R

Western blot analysis of IRAK-M in (M) m...



Immunohistochemistry of IRAK-M in human ...



Immunofluorescence of IRAK-M in Rat Live...