

---

## Product Datasheet

### LAG3 Antibody (orb1239549)

**Description**

LAG3 Antibody

**Species/Host**

Mouse

**Reactivity**

Human

**Conjugation**

Unconjugated

**Tested Applications**

ICC, IF, IHC-P

**Immunogen**

LAG-3 antibody was raised against the extracellular domain of human LAG-3

**Target**

LAG3

**Preservatives**

LAG-3 Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol.

**Form/Appearance**

Liquid

**Concentration**

1 mg/mL

**Storage**

LAG-3 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Note**

For research use only

**Application notes**

LAG-3 antibody can be used for Immunocytochemistry start at 1 µg/mL. For Immunofluorescence start at 10 µg/mL. For Immunohistochemistry start at 5 µg/mL. Antibody validated: Immunohistochemistry in human samples; Immunocytochemistry in human samples; Immunofluorescence in human samples and Flow Cytometry in human samples. All other applications and species not yet tested.

**Isotype**


IgG1,k

**Clonality**


Monoclonal

**Uniprot ID**
**P18627**
**NCBI**
**NP\_002277**
**Dilution Range**


LAG-3 antibody can be used for Immunocytochemistry start at 1 µg/mL. For Immunofluorescence start at 10 µg/mL. For Immunohistochemistry start at 5 µg/mL. Antibody validated: Immunohistochemistry in human samples; Immunocytochemistry in human samples; Immunofluorescence in human samples and Flow



Immunocytochemistry of LAG-3 in over exp...



Immunofluorescence of LAG-3 in over expr...



Immunofluorescence of LAG-3 in over huma...