



## **Product Datasheet**

BIRC8 Antibody (orb1239646)



Description nts. BIRC8 Antibody

Species/Host Rabbit

Reactivity Human, Mouse, Rat

**Conjugation** Unconjugated

Tested ELISA, ICC, IF, WB

**Applications** 

**Immunogen** ILP-2 antibody was raised with against a synthetic

peptide corresponding to amino acids near the amino terminus of human ILP-2. The immunogen is located

within the first 50 amino acids of ILP-2.

Target BIRC8

**Preservatives** ILP-2 Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/mL

**Storage** ILP-2 antibody can be stored at 4°C for three months

and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed

to prolonged high temperatures.

**Note** For research use only

**Application notes** ILP-2 antibody can be used for detection of ILP-2 by

Western blot at 1  $\mu$ g/mL. A band at approximately 33 kDa can be detected. Antibody can also be used for immunocytochemistry starting at 10  $\mu$ g/mL. For immunofluorescence start at 20  $\mu$ g/mL.Antibody validated: Western Blot in human samples; Immunocytochemistry in human samples and Immunofluorescence in human samples. All other

applications and species not yet tested.

**Isotype** IgG

**Clonality** Polyclonal

MW 33 kDa

Uniprot ID Q96P09


NCBI NP\_203127

**Dilution Range** ILP-2 antibody can be used for detection of ILP-2 by

Western blot at 1  $\mu$ g/mL. A band at approximately 33 kDa can be detected. Antibody can also be used for

immunocytochomistry starting at 10 ug/ml For


www.biorbyt.com



Western blot analysis of ILP-2 expressio...



Immunocytochemistry of ILP-2 in HepG2 ce...



Immunofluorescence of ILP-2 in HepG2 cel...