



Product Datasheet

Dffa Antibody (orb1239702)

Description	Dffa Antibody
Species/Host	Rabbit
Reactivity	Mouse
Conjugation	Unconjugated
Tested Applications	ELISA, IF, IHC-P, WB
Immunogen	ICAD antibody was raised against a peptide corresponding to amino acids near the amino terminus of mouse ICAD. The immunogen is located within the first 50 amino acids of ICAD.
Target	Dffa
Preservatives	ICAD Antibody is supplied in PBS containing 0.02% sodium azide.
Form/Appearance	Liquid
Concentration	1 mg/ml
Storage	ICAD antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Note	For research use only
Application notes	ICAD antibody can be used for detection of ICAD by Western blot at 1 µg/mL. A 45 kDa band can be detected. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 10 µg/mL. Antibody validated: Western Blot in mouse samples; Immunohistochemistry in mouse samples and Immunofluorescence in mouse samples. All other applications and species not yet tested.
Isotype	IgG
Clonality	Polyclonal
MW	45 kDa
Uniprot ID	O54786
NCBI	O54786
Dilution Range	ICAD antibody can be used for detection of ICAD by Western blot at 1 µg/mL. A 45 kDa band can be detected. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For



Immunohistochemistry of ICAD in mouse lu...



Immunofluorescence of ICAD in Mouse Lung...