



Product Datasheet

CCL11 Antibody (orb1239834)



Descriptionnts.

www.biorbyt.com

10 2.5 1

Species/Host Rabbit

Conjugation Unconjugated

Tested ELISA, ICC, IF, WB **Applications**

Immunogen Eotaxin antibody was raised in rabbits against a

CCL11 Antibody

peptide corresponding to amino acids near the carboxy terminus of human eotaxin. The immunogen is located within the last 50 amino acids of Eotaxin.

Target CCL11

Preservatives Eotaxin Antibody is supplied in PBS containing 0.02%

sodium azide.

Form/Appearance Liquid

Concentration 1 mg/ml

Storage Eotaxin antibody can be stored at 4°C for three

months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed

to prolonged high temperatures.

Note For research use only

Application notes Eotaxin antibody can be used for detection of eotaxin

by Western blot at 1 μ g/mL. The recombinant protein can be used as positive control and monomer and homodimer are detected at approximate 9 and 18

kDa. Antibody can also be used for

immunocytochemistry starting at 5 μ g/mL. For immunofluorescence start at 10 μ g/mL.Antibody validated: Western Blot in human samples; Immunocytochemistry in human samples and Immunofluorescence in human samples. All other

applications and species not yet tested.

Isotype IgG

Clonality Polyclonal


MW 9 - 18 kDa

Uniprot ID P51671


NCBI CAB07027

Dilution Range Eotaxin antibody can be used for detection of eotaxin


by Western blot at 1 $\mu g/\text{mL}.$ The recombinant protein can be used as positive control and monomer and



Western blot analysis of eotaxin in HeLa...



Immunocytochemistry of Eotaxin in 293 ce...



Immunofluorescence of Eotaxin in 293 cel...