

Product Datasheet

BFAR Antibody (orb1240114)

Description

BFAR Antibody

Species/Host

Rabbit

Reactivity

Human, Mouse, Rat

Conjugation

Unconjugated

Tested Applications

ELISA, IF, IHC-P, WB

Immunogen

BFAR antibody was raised against a 14 amino acid synthetic peptide near the carboxy terminus of human BFAR. The immunogen is located within amino acids 300 - 350 of BFAR.

Target

BFAR

Preservatives

BFAR Antibody is supplied in PBS containing 0.02% sodium azide.

Form/Appearance

Liquid

Concentration

1 mg/mL

Storage

BFAR antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Note

For research use only

Application notes

BFAR antibody can be used for detection of BFAR by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in mouse samples and Immunofluorescence in mouse samples. All other applications and species not yet tested.

Isotype


IgG

Clonality


Polyclonal

Uniprot ID
[Q9NZS9](#)
NCBI
[NP_057645](#)
Dilution Range


BFAR antibody can be used for detection of BFAR by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in mouse samples and



Western blot analysis of BFAR in human k...



Immunohistochemistry of BFAR in mouse ki...



Immunofluorescence of BFAR in mouse kidn...