


Product Datasheet

TIGIT Antibody (orb1240146)


Description	TIGIT Antibody
Species/Host	Mouse
Reactivity	Human
Conjugation	Unconjugated
Tested Applications	ELISA, FC, ICC, IF, IHC-P, WB
Immunogen	TIGIT antibody was raised against the extracellular domain of human TIGIT
Target	TIGIT
Preservatives	TIGIT Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol.
Form/Appearance	Liquid
Concentration	1 mg/mL
Storage	TIGIT antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Note	For research use only
Application notes	TIGIT antibody can be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 1 µg/mL. For flow cytometry at 1 µg/ml. For immunocytochemistry at 1 µg/mL. For Western blot at 1 µg/mL. Antibody validated: Western Blot in human samples; Immunohistochemistry in human samples; Immunocytochemistry in human samples; Immunofluorescence in human samples and Flow Cytometry in human samples. All other applications and species not yet tested.
Isotype	IgG1
Clonality	Monoclonal
MW	Predicted: 26 kDa Observed: 47 kDa
Uniprot ID	Q495A1
NCBI	NP_776160
Dilution Range	TIGIT antibody can be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 1 µg/mL. For flow cytometry at 1 µg/ml. For immunocytochemistry at 1



Western blot analysis of TIGIT in over e...



Immunocytochemistry of TIGIT in over exp...



Immunofluorescence of TIGIT in over expr...