




---

## Product Datasheet

### HAGH Antibody (orb1240672)

| Description                | HAGH Antibody   |    |
|----------------------------|---|---|
| <b>Species/Host</b>        | Rabbit  |   |
| <b>Reactivity</b>          | Human   |   |
| <b>Conjugation</b>         | Unconjugated  |   |
| <b>Tested Applications</b> | ELISA, WB   | Antibody used in WB on Human Liver at 0....   |
| <b>Immunogen</b>           | Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of human HAGH.   |   |
| <b>Target</b>              | HAGH  |   |
| <b>Preservatives</b>       | Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.   |   |
| <b>Form/Appearance</b>     | Liquid  |    |
| <b>Concentration</b>       | batch dependent   |   |
| <b>Storage</b>             | For short periods of storage (days) store at 4°C. For longer periods of storage, store HAGH antibody at -20°C. As with any antibody avoid repeat freeze-thaw cycles.  | Antibody used in WB on Hum. Fetal Heart ...   |
| <b>Note</b>                | For research use only   |   |
| <b>Application notes</b>   | HAGH antibody can be used for detection of HAGH by ELISA at 1:312500. HAGH antibody can be used for detection of HAGH by western blot at 1 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. |  |
| <b>Clonality</b>           | Polyclonal  |   |
| <b>MW</b>                  | 29 kDa  |   |
| <b>Uniprot ID</b>          | <a href="#">Q16775</a>  | Antibody used in WB on Hum. Fetal Lung a...   |
| <b>NCBI</b>                | <a href="#">NP_001035517</a>  |   |
| <b>Dilution Range</b>      | HAGH antibody can be used for detection of HAGH by ELISA at 1:312500. HAGH antibody can be used for detection of HAGH by western blot at 1 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. |   |
| <b>Expiration Date</b>     | 12 months from date of receipt.   |   |