

Product Datasheet

C9orf96 Antibody (orb1241043)

| | | |
|----------------------------|--|---|
| Description | C9orf96 Antibody |  |
| Species/Host | Rabbit | |
| Reactivity | Human | |
| Conjugation | Unconjugated | |
| Tested Applications | ELISA, WB | Antibody used in WB on Human HepG2 at 2.... |
| Immunogen | Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of human C9orf96. | |
| Target | C9orf96 | |
| Preservatives | Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. | |
| Form/Appearance | Liquid | |
| Concentration | batch dependent | |
| Storage | For short periods of storage (days) store at 4°C. For longer periods of storage, store C9orf96 antibody at -20°C. As with any antibody avoid repeat freeze-thaw cycles. | |
| Note | For research use only | |
| Application notes | C9orf96 antibody can be used for detection of C9orf96 by ELISA at 1:1562500. C9orf96 antibody can be used for detection of C9orf96 by western blot at 2.5 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. | |
| Clonality | Polyclonal | |
| MW | 76 kDa | |
| Uniprot ID | Q8NE28 | |
| NCBI | NP_714921 | |
| Dilution Range | C9orf96 antibody can be used for detection of C9orf96 by ELISA at 1:1562500. C9orf96 antibody can be used for detection of C9orf96 by western blot at 2.5 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. | |
| Expiration Date | 12 months from date of receipt. | |