

Product Datasheet

WDR12 Antibody (orb1241913)

| | | |
|----------------------------|---|---|
| Description | WDR12 Antibody |  |
| Species/Host | Rabbit | |
| Reactivity | Canine, Human, Mouse, Rat | <p>Antibody used in WB on Human Jurkat 5.0 ...</p> |
| Conjugation | Unconjugated | |
| Tested Applications | ELISA, WB | |
| Immunogen | Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of human WDR12. | |
| Target | WDR12 | |
| Preservatives | Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. | |
| Form/Appearance | Liquid | |
| Concentration | batch dependent | |
| Storage | For short periods of storage (days) store at 4°C. For longer periods of storage, store WDR12 antibody at -20°C. As with any antibody avoid repeat freeze-thaw cycles. | |
| Note | For research use only | |
| Application notes | WDR12 antibody can be used for detection of WDR12 by ELISA at 1:312500. WDR12 antibody can be used for detection of WDR12 by western blot at 5.0 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. | |
| Clonality | Polyclonal | |
| MW | 47 kDa | |
| Uniprot ID | Q9GZL7 | |
| NCBI | NP_060726 | |
| Dilution Range | WDR12 antibody can be used for detection of WDR12 by ELISA at 1:312500. WDR12 antibody can be used for detection of WDR12 by western blot at 5.0 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. | |
| Expiration Date | 12 months from date of receipt. | |