





---

## Product Datasheet

### FUS Antibody (orb1242221)

<b>Description</b>	FUS Antibody	
<b>Species/Host</b>	Rabbit	
<b>Reactivity</b>	Human, Mouse	
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	ELISA, IHC, WB	
<b>Immunogen</b>	Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of human FUS.	Antibody used in IHC on Human Breast at ...
<b>Target</b>	FUS	
<b>Preservatives</b>	Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.	
<b>Form/Appearance</b>	Liquid	
<b>Concentration</b>	batch dependent	
<b>Storage</b>	For short periods of storage (days) store at 4°C. For longer periods of storage, store FUS antibody at -20°C. As with any antibody avoid repeat freeze-thaw cycles.	Antibody used in IHC on Human Intestine ...
<b>Note</b>	For research use only	
<b>Application notes</b>	FUS antibody can be used for detection of FUS by ELISA at 1:312500. FUS antibody can be used for detection of FUS by western blot at 0.25 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000.	
<b>Clonality</b>	Polyclonal	
<b>MW</b>	29 kDa, 53 kDa	
<b>Uniprot ID</b>	<a href="#">P35637</a>	Antibody used in WB on mouse brain at 1:...
<b>NCBI</b>	<a href="#">NP_001010850</a>	
<b>Dilution Range</b>	FUS antibody can be used for detection of FUS by ELISA at 1:312500. FUS antibody can be used for detection of FUS by western blot at 0.25 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000.	
<b>Expiration Date</b>	12 months from date of receipt.	