




Product Datasheet

MEOX1 Antibody (orb1243756)

| | | |
|----------------------------|---|---|
| Description | MEOX1 Antibody |  |
| Species/Host | Rabbit |  |
| Reactivity | Human | |
| Conjugation | Unconjugated | |
| Tested Applications | ELISA, IHC, WB | Antibody used in IHC on Human Stomach. |
| Immunogen | Antibody produced in rabbits immunized with a synthetic peptide corresponding a region of human MEOX1. | |
| Target | MEOX1 |  |
| Preservatives | Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. | Antibody used in IHC on Human Liver. |
| Form/Appearance | Liquid | |
| Concentration | batch dependent | |
| Storage | For short periods of storage (days) store at 4°C. For longer periods of storage, store MEOX1 antibody at -20°C. As with any antibody avoid repeat freeze-thaw cycles. | |
| Note | For research use only | |
| Application notes | MEOX1 antibody can be used for detection of MEOX1 by ELISA at 1:312500. MEOX1 antibody can be used for detection of MEOX1 by western blot at 0.2-2.0 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. |  |
| Clonality | Polyclonal | |
| MW | 28 kDa, 20 kDa | Antibody used in WB on Human Jurkat 0.2-... |
| Uniprot ID | P50221 | |
| NCBI | NP_004518 | |
| Dilution Range | MEOX1 antibody can be used for detection of MEOX1 by ELISA at 1:312500. MEOX1 antibody can be used for detection of MEOX1 by western blot at 0.2-2.0 µg/mL, and HRP conjugated secondary antibody should be diluted 1:50,000 - 100,000. | |
| Expiration Date | 12 months from date of receipt. | |